ELSEVIER

Neuroscience Letters 324 (2002) 256–260

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 324

Abe, K., see Omori, N., 145
Aiso, S., see Tajima, H., 227
Akiyama, T., see Kitagawa, H., 232
Alvarez, C.E., Sutcliffe, J.G., Hypocretin is an early member of the incretin gene family, 169
Andrade, R., see Chapin, E.M., 1
Arévalo-Martín, A., see Molina-Holgado, E., 237
Ariumi, H., Takano, Y., Masumi, A., Takahashi, S., Hirabara, Y., Honda, K., Saito, R., Kamiya, H.-o., Roles of the central prostaglandin EP3 receptors in cardiovascular regulation in rats, 61
Arvanitis, D.A., see Zetterberg, H., 189

Babbs, A.J., see Irving, E.A., 53
Baethmann, A., see Stoffel, M., 201
Balm, P.H.M., see Höglund, E., 193
Bang, J.H., see Kim, C., 185
Bardenheuer, H., see Schönknecht, P., 83
Beyreuther, K., see Schönknecht, P., 83
Birbaumer, N., see Kübler, A., 205
Bissoon, N., see Boonstra, R., 173
Blennow, K., see Zetterberg, H., 189
Boonstra, R., Takagi, N., Bissoon, N., Vij, S., Gurd, J.W., Trapping-induced changes in expression of the *N*-methyl-D-aspartate receptor in the hippocampus of snowshoe hares, 173
Bouras, C., see Bozikas, V.P., 109
Bozikas, V.P., Kövari, E., Bouras, C., Karavatos, A., Neurofibrillary tangles in elderly patients with late onset schizophrenia, 109
Brill, T., see Kornhuber, M.E., 93

Campbell, C.A., see Irving, E.A., 53
Carson-Henderson, L., see Henderson, J.N., 77
Castillo, J., see Revilla, M., 29
Cervera, Á., see Revilla, M., 29
Chamorro, Á., see Revilla, M., 29
Chapin, E.M., Haj-Dahmane, S., Torres, G., Andrade, R., The 5-HT_4 receptor-induced depolarization in rat hippocampal neurons is mediated by cAMP but is independent of I_h, 1
Chartier-Harlin, M.-C., see Green, E.K., 113
Chen, C.-J., see Liao, S.-L., 133
Cohen, L.G., see Kübler, A., 205
Crook, J., see Henderson, J.N., 77
Crook, R., see Henderson, J.N., 77

Dávalos, A., see Revilla, M., 29
Déglise, P., Grünewald, B., Gauthier, M., The insecticide imidacloprid is a partial agonist of the nicotinic receptor of honeybee Kenyon cells [Neurosci. Lett. 321 (2002) 13–16], 86
de Lima, A.D., see Wilhelmi, E., 255
Deininger, M.H., Trautmann, K., Magdolen, V., Luther, T., Schluesener, H.J., Meyermann, R., Cortical neurons of Creutzfeldt–Jakob disease patients express the urokinase-type plasminogen activator receptor, 80
Del Arco, A., see Díaz-Cabiale, Z., 154
Díaz-Cabiale, Z., Vivó, M., Del Arco, A., O'Connor, W.T., Harte, M.K., Müller, C.E., Martínez, E., Popoli, P., Fuxe, K., Ferré, S., Metabotropic glutamate mGlu5 receptor-mediated modulation of the ventral striopallidal GABA pathway in rats. Interactions with adenosine A_{2A} and dopamine D_2 receptors, 154
Diez-Martínez, O., see Manjarrez, E., 213
Diggle, P., see Pendleton, N., 74
Doppelmayr, M., see Sauseng, P., 121

Eaton, M.J., see Hains, B.C., 222
Eder, E., see Götz, M.E., 49
Eisner, W., see Golaszewski, S.M., 125
Engelmann, M., Wolf, G., Horn, T.F.W., Release patterns of excitatory and inhibitory amino acids within the hypothalamic supraoptic nucleus in response to direct nitric oxide administration during forced swimming in rats, 252

Fekete, C., see Wittmann, G., 69
Felber, S., see Golaszewski, S.M., 125
Ferré, S., see Díaz-Cabiale, Z., 154
Flajolet, M., see Leypoldt, F., 37
Flonta, M.-L., see Reid, G., 164
Flores, A., see Manjarrez, E., 213
Fujii, S., see Yamazaki, Y., 65
Fukui, K., see Urai, Y., 101
Funahashi, H., see Suzuki, R., 5
Fuxe, K., see Díaz-Cabiale, Z., 154

Ganz, C., see Kornhuber, M.E., 93
Gauthier, M., see Déglise, P., 86
Gérard, H., see Jean-Charles, V., 149
Ghribi, O., Herman, M.M., Savory, J., The endoplasmic reticulum is the main site for caspase-3 activation following aluminum-induced neurotoxicity in rabbit hippocampus, 217
Golaszewski, S.M., Zschiegner, F., Siedentopf, C.M., Unterrainer, J., Sweeney, R.A., Eisner, W., Lechner-Steinleitner, S., Mottaghy, F.M., Felber, S., A new pneumatic vibrator for functional magnetic resonance imaging of the human sensorimotor cortex, 125
Götz, M.E., Wacker, M., Luckhaus, C., Wanek, P., Tatschner, T., Jellinger, K., Leblhuber, F., Ransmayr, G., Riederer, P., Eder, E., Unaltered brain levels of 1,N^2-propanodeoxyguanosine adducts of *trans*-4-hydroxy-2-nonenal in Alzheimer's disease, 49
Graf, R., see Stoffel, M., 201

PII: S0304-3940(02)00434-2

Grass, S., Xu, I.S., Wiesenfeld-Hallin, Z., Xu, X.-J., Comparison of the effect of intrathecal endomorphin-1 and endomorphin-2 on spinal cord excitability in rats, 197
Green, A.J.E., Ramljak, S., Müller, W.E.G., Knight, R.S.G., Schröder, H.C., 14-3-3 in the cerebrospinal fluid of patients with variant and sporadic Creutzfeldt–Jakob disease measured using capture assay able to detect low levels of 14-3-3 protein, 57
Green, E.K., Thaker, U., McDonagh, A.M., Iwatsubo, T., Lambert, J.-C., Chartier-Harlin, M.-C., Harris, J.M., Pickering-Brown, S.M., Lendon, C.L., Mann, D.M.A., A polymorphism within intron 11 of the tau gene is not increased in frequency in patients with sporadic Alzheimer's disease, nor does it influence the extent of tau pathology in the brain, 113
Gruber, W., see Sauseng, P., 121
Grünewald, B., see Déglise, P., 86
Gu, Y., Huang, L.-Y.M., Gabapentin potentiates *N*-methyl-D-aspartate receptor mediated currents in rat GABAergic dorsal horn neurons, 177
Guan, J.-L., see Suzuki, R., 5
Guaza, C., see Molina-Holgado, E., 237
Guo, H., see Tang, Z., 13
Gurd, J.W., see Boonstra, R., 173

Habara, Y., see Takahashi-Iwanaga, H., 137
Hahm, D.-H., see Yun, S.-J., 45
Hains, B.C., Yucra, J.A., Eaton, M.J., Hulsebosch, C.E., Intralesion transplantation of serotonergic precursors enhances locomotor recovery but has no effect on development of chronic central pain following hemisection injury in rats, 222
Hajak, G., see Rodenbeck, A., 159
Haj-Dahmane, S., see Chapin, E.M., 1
Hardy, J., see Henderson, J.N., 77
Harris, J.M., see Green, E.K., 113
Harrison, D.C., see Irving, E.A., 53
Harte, M.K., see Díaz-Cabiale, Z., 154
Hartmann, T., see Schönknecht, P., 83
Hashimoto, Y., see Tajima, H., 227
Hayakawa, T., see Omori, N., 145
Henderson, J.N., Crook, R., Crook, J., Hardy, J., Onstead, L., Carson-Henderson, L., Mayer, P., Parker, B., Petersen, R., Williams, B., Apolipoprotein E4 and tau allele frequencies among Choctaw Indians, 77
Henshall, D.C., see Meller, R., 33
Herman, M.M., see Ghribi, O., 217
Hinterberger, T., see Kübler, A., 205
Hirabara, Y., see Ariumi, H., 61
Höglund, E., Balm, P.H.M., Winberg, S., Stimulatory and inhibitory effects of 5-HT_{1A} receptors on adrenocorticotropic hormone and cortisol secretion in a teleost fish, the Arctic charr (*Salvelinus alpinus*), 193
Holland, F., see Pendleton, N., 74
Honda, K., see Ariumi, H., 61
Horan, M.A., see Pendleton, N., 74
Horn, T.F.W., see Engelmann, M., 252
Hu, G.-Y., see Li, Y., 25
Huang, L.-Y.M., see Gu, Y., 177
Huether, G., see Rodenbeck, A., 159
Hulsebosch, C.E., see Hains, B.C., 222
Hunter, A.J., see Irving, E.A., 53

Ioka, M., see Narita, M., 97
Irving, E.A., Harrison, D.C., Babbs, A.J., Mayes, A.C., Campbell, C.A., Hunter, A.J., Upton, N., Parsons, A.A., Increased cortical expression of the orexin-1 receptor following permanent middle cerebral artery occlusion in the rat, 53
Ito, K.-I., see Yamazaki, Y., 65
Ito, Y., see Tajima, H., 227
Iwatsubo, T., see Green, E.K., 113

Jang, C.H., see Kim, C., 185
Jean-Charles, V., Gérard, H., Noradrenergic receptors and in vitro respiratory rhythm: possible interspecies differences between mouse and rat neonates, 149
Jellinger, K., see Götz, M.E., 49
Jiang, X.-F., see Namura, S., 117
Jinnouchi, O., see Urai, Y., 101
Joo, I., see Kim, C., 185
Jung, M.W., see Kim, C., 185

Kamichi, S., see Namura, S., 117
Kamiya, H.-o., see Ariumi, H., 61
Kaneko, T., see Li, J.-L., 9
Karavatos, A., see Bozikas, V.P., 109
Kariya, K., see Tanaka, J., 242
Kato, H., see Yamazaki, Y., 65
Kawada, T., see Kitagawa, H., 232
Kawatani, M., see Sato, M., 89
Kiening, K.L., van Landeghem, F.K.H., Schreiber, S., Thomale, U.W., von Deimling, A., Unterberg, A.W., Stover, J.F., Decreased hemispheric Aquaporin-4 is linked to evolving brain edema following controlled cortical impact injury in rats, 105
Kikuyama, S., see Suzuki, R., 5
Kılıcturgay, S., see Özarda İlçöl, Y., 41
Kim, C., Jang, C.H., Bang, J.H., Jung, M.W., Joo, I., Kim, S.U., Mook-Jung, I., Amyloid precursor protein processing is separately regulated by protein kinase C and tyrosine kinase in human astrocytes, 185
Kim, S.U., see Kim, C., 185
Kita, Y., see Tajima, H., 227
Kitagawa, H., Yamazaki, T., Akiyama, T., Yahagi, N., Kawada, T., Mori, H., Sunagawa, K., Modulatory effects of ketamine on catecholamine efflux from in vivo cardiac sympathetic nerve endings in cats, 232
Klimesch, W., see Sauseng, P., 121
Knight, R.S.G., see Green, A.J.E., 57
Kohsaka, S., see Omori, N., 145
Kornhuber, M.E., Ganz, C., Lang, R., Brill, T., Schmahl, W., Focal encephalitis in the Lewis rat induced by intracerebral enterotoxin superantigen and amplified by activated intravenous splenocytes, 93
Koto, A., see Tajima, H., 227
Kövari, E., see Bozikas, V.P., 109
Kübler, A., Schmidt, K., Cohen, L.G., Lotze, M., Winter, S., Hinterberger, T., Birbaumer, N., Modulation of slow cortical potentials by transcranial magnetic stimulation in humans, 205
Kudo, Y., see Yamazaki, Y., 65
Kwak, K.T., see Urai, Y., 101

Lambert, J.-C., see Green, E.K., 113
Lang, R., see Kornhuber, M.E., 93
Leblhuber, F., see Götz, M.E., 49
Lechan, R.M., see Wittmann, G., 69
Lechner-Steinleitner, S., see Golaszewski, S.M., 125
Lee, E.H., see Yun, S.-J., 45
Lendon, C.L., see Green, E.K., 113
Leypoldt, F., Flajolet, M., Methner, A., Neuronal differentiation of cultured human NTERA-2cl.D1 cells leads to increased expression of synapsins, 37

Li, J.-L., Kaneko, T., Mizuno, N., Synaptic association of dopaminergic axon terminals and neurokinin-1 receptor-expressing intrinsic neurons in the striatum of the rat, 9
Li, Y., Hu, G.-Y., Huperzine A, a nootropic agent, inhibits fast transient potassium current in rat dissociated hippocampal neurons, 25
Liang, C., see Wei, H., 181
Liao, S.-L., Raung, S.-L., Chen, C.-J., Japanese encephalitis virus stimulates superoxide dismutase activity in rat glial cultures, 133
Linke, R., see Wilhelmi, E., 255
Liposits, Z., see Wittmann, G., 69
Lotze, M., see Kübler, A., 205
Lubec, G., see Shim, K.S., 209
Luckhaus, C., see Götz, M.E., 49
Luther, T., see Deininger, M.H., 80
Lütjohann, D., see Schönknecht, P., 83

Maeno, H., see Namura, S., 117
Magdolen, V., see Deininger, M.H., 80
Manjarrez, E., Diez-Martínez, O., Méndez, I., Flores, A., Stochastic resonance in human electroencephalographic activity elicited by mechanical tactile stimuli, 213
Mann, D.M.A., see Green, E.K., 113
Martínez, E., see Díaz-Cabiale, Z., 154
Masumi, A., see Ariumi, H., 61
Mayer, P., see Henderson, J.N., 77
Mayes, A.C., see Irving, E.A., 53
McDonagh, A.M., see Green, E.K., 113
McDowell, K., The effects of probability and relative direction on human somatomotor electroencephalographic rhythms, 17
Meller, R., Skradski, S.L., Simon, R.P., Henshall, D.C., Expression, proteolysis and activation of caspases 6 and 7 during rat C6 glioma cell apoptosis, 33
Méndez, I., see Manjarrez, E., 213
Methner, A., see Leypoldt, F., 37
Meyermann, R., see Deininger, M.H., 80
Miyakawa, H., see Yamazaki, Y., 65
Miyakubo, H., see Tanaka, J., 242
Mizuguchi, H., see Omori, N., 145
Mizuno, N., see Li, J.-L., 9
Molina-Holgado, E., Arévalo-Martín, A., Ortiz, S., Vela, J.M., Guaza, C., Theiler's virus infection induces the expression of cyclooxygenase-2 in murine astrocytes: inhibition by the anti-inflammatory cytokines interleukin-4 and interleukin-10, 237
Mook-Jung, I., see Kim, C., 185
Mori, H., see Kitagawa, H., 232
Mottaghy, F.M., see Golaszewski, S.M., 125
Müller, C.E., see Díaz-Cabiale, Z., 154
Müller, W.E.G., see Green, A.J.E., 57

Nagahiro, S., see Urai, Y., 101
Nagata, I., see Namura, S., 117
Nakajo, S., see Suzuki, R., 5
Nakamura, T., see Yamazaki, Y., 65
Nakazawa, K., Ojima, H., Ohno, Y., A highly conserved tryptophane residue indispensable for cloned rat neuronal P2X receptor activation, 141
Namura, S., Maeno, H., Takami, S., Jiang, X.-F., Kamichi, S., Wada, K., Nagata, I., Inhibition of glial glutamate transporter GLT-1 augments brain edema after transient focal cerebral ischemia in mice, 117
Narita, M., see Narita, M., 97
Narita, M., Ioka, M., Suzuki, M., Narita, M., Suzuki, T., Effect of repeated administration of morphine on the activity of extracellular signal regulated kinase in the mouse brain, 97
Narita, M., see Narita, M., 129
Narita, M., Soma, M., Tamaki, H., Narita, M., Suzuki, T., Intensification of the development of ethanol dependence in mice lacking dopamine D_3 receptor, 129
Niikura, T., see Tajima, H., 227
Nishimoto, I., see Tajima, H., 227
Nomura, M., see Tanaka, J., 242

Obach, V., see Revilla, M., 29
O'Connor, W.T., see Díaz-Cabiale, Z., 154
Ohno, Y., see Nakazawa, K., 141
Ohsawa, K., see Omori, N., 145
Ojima, H., see Nakazawa, K., 141
Ollier, W.E.R., see Pendleton, N., 74
Omori, N., Mizuguchi, H., Ohsawa, K., Kohsaka, S., Hayakawa, T., Abe, K., Shibasaki, F., Modification of a fiber protein in an adenovirus vector improves in vitro gene transfer efficiency to the mouse microglial cell line, 145
Onstead, L., see Henderson, J.N., 77
Ortiz, S., see Molina-Holgado, E., 237
Özarda İlçöl, Y., Özyurt, G., Kılıcturgay, S., Uncu, G., Ulus, İ.H., The decline in serum choline concentration in humans during and after surgery is associated with the elevation of cortisol, adrenocorticotropic hormone, prolactin and β-endorphin concentrations, 41
Özyurt, G., see Özarda İlçöl, Y., 41

Palmér, M., see Zetterberg, H., 189
Palmqvist, L., see Zetterberg, H., 189
Pantel, J., see Schönknecht, P., 83
Pape, H.-C., see Wilhelmi, E., 255
Parker, B., see Henderson, J.N., 77
Parsons, A.A., see Irving, E.A., 53
Payton, A., see Pendleton, N., 74
Pendleton, N., Payton, A., van den Boogerd, E.H., Holland, F., Diggle, P., Rabbitt, P.M.A., Horan, M.A., Worthington, J., Ollier, W.E.R., Apolipoprotein E genotype does not predict decline in intelligence in healthy older adults, 74
Petersen, R., see Henderson, J.N., 77
Pickering-Brown, S.M., see Green, E.K., 113
Plesnila, N., see Stoffel, M., 201
Poirier, J., see Zetterberg, H., 189
Popoli, P., see Díaz-Cabiale, Z., 154

Rabbitt, P.M.A., see Pendleton, N., 74
Ramljak, S., see Green, A.J.E., 57
Ransmayr, G., see Götz, M.E., 49
Raung, S.-L., see Liao, S.-L., 133
Reid, G., Flonta, M.-L., Ion channels activated by cold and menthol in cultured rat dorsal root ganglion neurones, 164
Revilla, M., Obach, V., Cervera, Á., Dávalos, A., Castillo, J., Chamorro, Á., A -174G/C polymorphism of the interleukin-6 gene in patients with lacunar infarction, 29
Ricksten, A., see Zetterberg, H., 189
Riederer, P., see Götz, M.E., 49
Rinecker, M., see Stoffel, M., 201
Rodenbeck, A., Huether, G., Rüther, E., Hajak, G., Interactions between evening and nocturnal cortisol secretion and sleep parameters in patients with severe chronic primary insomnia, 159

Rüther, E., see Rodenbeck, A., 159
Rymo, L., see Zetterberg, H., 189

Saito, R., see Ariumi, H., 61
Sakamaki, K., see Tanaka, J., 242
Sato, M., Kawatani, M., Effects of noradrenaline on cytosolic concentrations of Ca^{2+} in cultured corpus cavernosum smooth muscle cells of the rabbit, 89
Sauseng, P., Klimesch, W., Gruber, W., Doppelmayr, M., Stadler, W., Schabus, M., The interplay between theta and alpha oscillations in the human electroencephalogram reflects the transfer of information between memory systems, 121
Savory, J., see Ghribi, O., 217
Schabus, M., see Sauseng, P., 121
Schluesener, H.J., see Deininger, M.H., 80
Schmahl, W., see Kornhuber, M.E., 93
Schmidt, K., see Kübler, A., 205
Schönknecht, P., Lütjohann, D., Pantel, J., Bardenheuer, H., Hartmann, T., von Bergmann, K., Beyreuther, K., Schröder, J., Cerebrospinal fluid 24S-hydroxycholesterol is increased in patients with Alzheimer's disease compared to healthy controls, 83
Schreiber, S., see Kiening, K.L., 105
Schröder, H.C., see Green, A.J.E., 57
Schröder, J., see Schönknecht, P., 83
Shibasaki, F., see Omori, N., 145
Shim, K.S., Lubec, G., Drebrin, a dendritic spine protein, is manifold decreased in brains of patients with Alzheimer's disease and Down syndrome, 209
Shimojima, H., see Suzuki, R., 5
Shioda, S., see Suzuki, R., 5
Shou, T., see Zhan, X., 247
Siedentopf, C.M., see Golaszewski, S.M., 125
Simon, R.P., see Meller, R., 33
Skradski, S.L., see Meller, R., 33
Soma, M., see Narita, M., 129
Spandidos, D.A., see Zetterberg, H., 189
Stadler, W., see Sauseng, P., 121
Stoffel, M., Rinecker, M., Graf, R., Baethmann, A., Plesnila, N., Nitric oxide in the penumbra of a focal cortical necrosis in rats, 201
Stover, J.F., see Kiening, K.L., 105
Sunagawa, K., see Kitagawa, H., 232
Sutcliffe, J.G., see Alvarez, C.E., 169
Suzue, A., see Urai, Y., 101
Suzuki, H., Yamamoto, T., Centrifugal neurons of the octopus optic lobe cortex are immunopositive for calcitonin gene-related peptide, 21
Suzuki, M., see Narita, M., 97
Suzuki, R., Shimojima, H., Funahashi, H., Nakajo, S., Yamada, S., Guan, J.-L., Tsurugano, S., Uehara, K., Takeyama, Y., Kikuyama, S., Shioda, S., Orexin-1 receptor immunoreactivity in chemically identified target neurons in the rat hypothalamus, 5
Suzuki, T., see Narita, M., 97
Suzuki, T., see Narita, M., 129
Sweeney, R.A., see Golaszewski, S.M., 125

Tajima, H., Niikura, T., Hashimoto, Y., Ito, Y., Kita, Y., Terashita, K., Yamazaki, K., Koto, A., Aiso, S., Nishimoto, I., Evidence for in vivo production of Humanin peptide, a neuroprotective factor against Alzheimer's disease-related insults, 227
Takagi, N., see Boonstra, R., 173
Takahashi, S., see Ariumi, H., 61
Takahashi-Iwanaga, H., Habara, Y., Adenosine 5′-triphosphate-evoked calcium responses in terminal Schwann cells of lanceolate sensory endings isolated from rat vibrissae, 137
Takami, S., see Namura, S., 117
Takano, Y., see Ariumi, H., 61
Takeyama, Y., see Suzuki, R., 5
Tamaki, H., see Narita, M., 129
Tanaka, J., Kariya, K., Miyakubo, H., Sakamaki, K., Nomura, M., Attenuated drinking response induced by angiotensinergic activation of subfornical organ projections to the paraventricular nucleus in estrogen-treated rats, 242
Tang, Z., Yu, Y., Guo, H., Zhou, J., Induction of tyrosine hydroxylase expression in rat fetal striatal precursor cells following transplantation, 13
Tatschner, T., see Götz, M.E., 49
Terashita, K., see Tajima, H., 227
Thaker, U., see Green, E.K., 113
Thomale, U.W., see Kiening, K.L., 105
Torres, G., see Chapin, E.M., 1
Trautmann, K., see Deininger, M.H., 80
Tsurugano, S., see Suzuki, R., 5

Uehara, K., see Suzuki, R., 5
Ulus, İ.H., see Özarda İlçöl, Y., 41
Uncu, G., see Özarda İlçöl, Y., 41
Unterberg, A.W., see Kiening, K.L., 105
Unterrainer, J., see Golaszewski, S.M., 125
Upton, N., see Irving, E.A., 53
Urai, Y., Jinnouchi, O., Kwak, K.T., Suzue, A., Nagahiro, S., Fukui, K., Gene expression of D-amino acid oxidase in cultured rat astrocytes: regional and cell type specific expression, 101

van den Boogerd, E.H., see Pendleton, N., 74
van Landeghem, F.K.H., see Kiening, K.L., 105
Vela, J.M., see Molina-Holgado, E., 237
Vij, S., see Boonstra, R., 173
Vivó, M., see Díaz-Cabiale, Z., 154
von Bergmann, K., see Schönknecht, P., 83
von Deimling, A., see Kiening, K.L., 105

Wacker, M., see Götz, M.E., 49
Wada, K., see Namura, S., 117
Wanek, P., see Götz, M.E., 49
Wei, H., Xiong, W., Yang, S., Zhou, Q., Liang, C., Zeng, B.X., Xu, L., Propofol facilitates the development of long-term depression (LTD) and impairs the maintenance of long-term potentiation (LTP) in the CA1 region of the hippocampus of anesthetized rats, 181
Wiesenfeld-Hallin, Z., see Grass, S., 197
Wilhelmi, E., Linke, R., de Lima, A.D., Pape, H.-C., Erratum to ''Axonal connections of thalamic posterior paralaminar nuclei with amygdaloid projection neurons to the cholinergic basal forebrain in the rat'' [Neurosci. Lett. 315 (2001) 121–124], 255
Williams, B., see Henderson, J.N., 77
Winberg, S., see Höglund, E., 193
Winter, S., see Kübler, A., 205
Wittmann, G., Liposits, Z., Lechan, R.M., Fekete, C., Medullary adrenergic neurons contribute to the neuropeptide Y-ergic innervation of hypophysiotropic thyrotropin-releasing hormone-synthesizing neurons in the rat, 69
Wolf, G., see Engelmann, M., 252
Worthington, J., see Pendleton, N., 74

Xiong, W., see Wei, H., 181
Xu, I.S., see Grass, S., 197
Xu, L., see Wei, H., 181
Xu, X.-J., see Grass, S., 197

Yahagi, N., see Kitagawa, H., 232
Yamada, S., see Suzuki, R., 5
Yamamoto, T., see Suzuki, H., 21
Yamazaki, K., see Tajima, H., 227
Yamazaki, T., see Kitagawa, H., 232
Yamazaki, Y., Fujii, S., Nakamura, T., Miyakawa, H., Kudo, Y., Kato, H., Ito, K.-I., Changes in $[Ca^{2+}]_i$ during adenosine triphosphate-induced synaptic plasticity in hippocampal CA1 neurons of the guinea pig, 65
Yang, S., see Wei, H., 181
Yu, Y., see Tang, Z., 13
Yucra, J.A., see Hains, B.C., 222
Yun, S.-J., Hahm, D.-H., Lee, E.H., Immobilization stress induces the expression of αB-crystallin in rat hippocampus: implications of glial activation in stress-mediated hippocampal degeneration, 45

Zafiropoulos, A., see Zetterberg, H., 189
Zeng, B.X., see Wei, H., 181
Zetterberg, H., Palmér, M., Ricksten, A., Poirier, J., Palmqvist, L., Rymo, L., Zafiropoulos, A., Arvanitis, D.A., Spandidos, D.A., Blennow, K., Influence of the apolipoprotein E ε4 allele on human embryonic development, 189
Zhan, X., Shou, T., Anatomical evidence of subcortical contributions to the orientation selectivity and columns of the cat's primary visual cortex, 247
Zhou, J., see Tang, Z., 13
Zhou, Q., see Wei, H., 181
Zschiegner, F., see Golaszewski, S.M., 125